

U.S. Serial No. 10/734,218
Page 2

Amendment and Response to Restriction Requirement

IN THE CLAIMS

Please amend the claims to read as follows:

1.-25. (Cancelled)

26. (Currently Amended) A polymerization catalyst obtained by a process that comprises
reacting:

an aluminum oxy compound(a) having a ratio of an intensity at 30 ppm(H2) to an intensity
at 10 ppm(H1) [H2/H1] in an ²⁷Al-solid NMR spectrum of less than 0.35,

water(b), and

a compound(c) having a hydroxy group selected from the group consisting of tertiary
alcohols and alcohols substitute with a halogen atom, the alcohols substituted with a halogen being
selected from the group consisting of alcohols indicated by the formula CR¹R²R³-OH (wherein each
of R¹, R² and R³ independently represents a hydrogen atom or hydrocarbon group having 1 to 20
carbon atoms, which is optionally substituted with a halogen atom, and they may be mutually the
same or different), halogenated phenol compounds and silanol compounds;

wherein the modified aluminum oxy compound (A) has a ratio of an intensity at 30 ppm(M2)
to an intensity at 10 ppm(M1)[M2/M1] in an ²⁷Al-solid NMR spectrum of 0.60 or more, the molar
ratio[(a)/(b)] of the aluminum oxy compound(a) to water(b) is 1/3 to 1/0.01, and the molar ratio
[(a)/(c)] of the aluminum oxy compound(a) to the compound(c) is 1/3 to 1/0.01 contacting (A) a
~~modified aluminum oxy compound, which is an aluminum oxy compound that has been modified
by reacting with a compound having a hydroxy group, wherein the modified aluminum oxy~~

U.S. Serial No. 10/734,218
Page 3

Amendment and Response to Restriction Requirement

~~compound has a ratio of an intensity at 30 ppm(M2) to an intensity at 10 ppm(M1) [M2/M1], in an ²⁷Al solid NMR spectrum of 0.60 or more, and contacting said modified aluminum oxy compound(A) with (B) a transition metal compound.~~

27. (Previously presented) A polymerization catalyst obtained by a process that comprises contacting (A) a modified aluminum oxy compound, obtained by a process that comprises reacting:

an aluminum oxy compound (a) having a ratio of an intensity at 30 ppm(H2) to an intensity at 10 ppm(H1) [H2/H1] in an ²⁷Al-solid NMR spectrum of less than 0.35,

water (b), and

a compound (c) having a hydroxy group, wherein the modified aluminum oxy compound has a ratio of an intensity at 30 ppm(M2) to an intensity at 30 ppm(M1) [M2/M1] in an ²⁷Al-solid NMR spectrum of 0.60 or more,

with (B) a transition metal compound.

28. (Currently amended) A polymerization catalyst obtained by a process that comprises reacting:

an aluminum oxy compound(a') having a ratio of an intensity at 30 ppm(L2) to an intensity at 10 ppm(L1) [L2/L1] in an ²⁷Al-solid NMR spectrum of not less than 0.35, which is obtained by reacting an aluminum oxy compound(a) having a ratio of an intensity at 30 ppm(H2) to an intensity at 10 ppm(H1)[H2/H1] in an ²⁷Al-solid NMR spectrum of less than 0.35 and water (b); and

U.S. Serial No. 10/734,218
Page 4

Amendment and Response to Restriction Requirement

a compound(c) having a hydroxy group selected from the group consisting of tertiary alcohols and alcohols substituted with a halogen atom, the alcohols substituted with a halogen being selected from alcohols indicated by the formula $CR^1R^2R^3-OH$ (wherein each of R^1 , R^2 and R^3 independently represents a hydrogen atom or hydrocarbon group having 1 to 20 carbon atoms, which is optionally substituted with a halogen atom, and they may be mutually the same or different), halogenated phenol compounds and silanol compounds, wherein the modified aluminum oxy compound (A) has a ratio of an intensity at 30 ppm(M2) to an intensity at 10 ppm(ml) $[M2/M1]$ in an ^{27}Al -solid NMR spectrum of 0.60 or more, the molar ratio[(a)/(b)] of the aluminum oxy compound(a) to water(b) is 1/3 to 1/0.01, and the molar ratio[(a)/(c)] of the aluminum oxy compound(a) to the compound(c) is 1/3 to 1/0.01 and contacting said aluminum oxy compound (A)

with (B) a transition metal compound.

29. (Previously presented) A polymerization catalyst obtained by a process that comprises contacting (A) a modified aluminum oxy compound, obtained by a process that comprises reacting:

an aluminum oxy compound (a') having a ratio of an intensity at 30 ppm(L2) to an intensity at 10 ppm(L1) $[L2/L1]$ in an ^{27}Al -solid NMR spectrum of not less than 0.35, which is obtained by reacting an aluminum oxy compound (a) having a ratio of an intensity at 30 ppm(H2) to an intensity at 10 ppm (H1) $[H2/H1]$ in an ^{27}Al -solid NMR spectrum of less than 0.35 and water (b); and

a compound (c) having a hydroxy group

U.S. Serial No. 10/734,218
Page 5

Amendment and Response to Restriction Requirement

and (B) a transition metal compound.

30. (Currently Amended) A polymerization catalyst obtained by a process that comprises reacting:

an aluminum oxy compound(a'') having a ratio of an intensity at 30 ppm(N2) to an intensity at 10 ppm(N1) [N2/N1] in an ²⁷Al-solid NMR spectrum of not less than 0.35 obtained by reacting an aluminum oxy compound(a) having a ratio of an intensity at 30 ppm(H2) to an intensity at 10 ppm H1[H2/H1] in its ²⁷Al-solid NMR spectrum of less than 0.35 and a compound(c) having a hydroxy group selected from the group consisting of tertiary alcohols and alcohols substituted with a halogen atom, the alcohol being selected from alcohols indicated by the formula CR¹R²R³-OH (wherein each of R¹, R² and R³ independently represents a hydrogen atom or hydrocarbon group having 1 to 20 carbon atoms, which is optionally substituted with a halogen atom, and they may be mutually the same or different), halogenated phenol compounds and silanol compounds, with

water(b), wherein the modified aluminum oxy compound (A) has a ratio of an intensity at 30 ppm(M2) to an intensity at 10 ppm(M1)[M2/M1] in an ²⁷Al-solid NMR spectrum of 0.60 or more, the molar ratio[(a)/(b)] of the aluminum oxy compound(a) to water(b) is 1/3 to 1/0.01, and the molar ratio[(a)/(c)] of the aluminum oxy compound(a) to the compound(c) is 1/3 to 1/0.01 ; and
~~contacting (A) a modified aluminum oxy compound, obtained by a process that comprises reacting:~~
~~—— an aluminum oxy compound (a'') having a ratio of an intensity at 30 ppm(N2) to an intensity at 10 ppm(N1) [N2/N1] in an ²⁷Al-solid NMR spectrum of not less than 0.35 obtained by reacting an aluminum oxy compound (a) having a ratio of an intensity at 30 ppm(L2) to an intensity at 10~~

U.S. Serial No. 10/734,218
Page 6

Amendment and Response to Restriction Requirement

~~ppm (L1) [L2/L1] in its ²⁷Al solid NMR spectrum of less than 0.35 and a compound (c) having a hydroxy group, and~~

~~— water (c)~~

and (B) a transition metal compound.

31. (Previously presented) A polymerization catalyst obtained by a process that comprises contacting either:

(A) the modified aluminum oxy compound of claim 26,

(B) a transition metal compound, and

(C) an organoaluminum compound; or

(A) the modified aluminum oxy compound of claim 26,

(B) a transition metal compound,

(C) an organoaluminum compound, and

(D) any one of (D1) a boron compound represented by the general formula $BQ^1Q^2Q^3$, (D2) a boron compound represented by the general formula $G^+(BQ^1Q^2Q^3Q^4)^-$ and (D3) a boron compound represented by the general formula $(L-H)^+(BQ^1Q^2Q^3Q^4)^-$, wherein B represents a boron atom in the trivalent valence state; Q^1 to Q^3 may be the same or different and represent a halogen atom, a hydrocarbon group, a halogenated hydrocarbon group, a substituted silyl group, an alkoxy group or a di-substituted amino group; G^+ represents an inorganic or organic cation; and $(L-H)^+$ represents a Brønsted acid.

U.S. Serial No. 10/734,218
Page 7

Amendment and Response to Restriction Requirement

32. (Currently amended) A polymerization catalyst obtained by a process that comprises contacting either:

(A) the modified aluminum oxy compound of ~~claim~~ claim 27,

(B) a transition metal compound, and

(C) an organoaluminum compound; or

(A) the modified aluminum oxy compound of ~~claim~~ claim 27,

(B) a transition metal compound,

(C) an organoaluminum compound, and

(D) any one of (D1) a boron compound represented by the general formula $BQ^1Q^2Q^3$, (D2) a boron compound represented by the general formula $G^+(BQ^1Q^2Q^3Q^4)^-$ and (D3) a boron compound represented by the general formula $(L-H)^+(BQ^1Q^2Q^3Q^4)^-$, wherein B represents a boron atom in the trivalent valence state; Q^1 to Q^3 may be the same or different and represent a halogen atom, a hydrocarbon group, a halogenated hydrocarbon group, a substituted silyl group, an alkoxy group or a di-substituted amino group; G^+ represents an inorganic or organic cation; and $(L-H)^+$ represents a Brønsted acid.

33. (Previously presented) A polymerization catalyst obtained by a process that comprises contacting either:

(A) the modified aluminum oxy compound of claim 28,

U.S. Serial No. 10/734,218
Page 8

Amendment and Response to Restriction Requirement

(B) a transition metal compound, and

(C) an organoaluminum compound; or

(A) the modified aluminum oxy compound of claim 28,

(B) a transition metal compound,

(C) an organoaluminum compound, and

(D) any one of (D1) a boron compound represented by the general formula $BQ^1Q^2Q^3$, (D2) a boron compound represented by the general formula $G^+(BQ^1Q^2Q^3Q^4)^-$ and (D3) a boron compound represented by the general formula $(L-H)^+(BQ^1Q^2Q^3Q^4)^-$; wherein B represents a boron atom in the trivalent valence state; Q^1 to Q^3 may be the same or different and represent a halogen atom, a hydrocarbon group, a halogenated hydrocarbon group, a substituted silyl group, an alkoxy group or a di-substituted amino group; G^+ represents an inorganic or organic cation; and $(L-H)^+$ represents a Brønsted acid.

34. (Previously presented) A polymerization catalyst obtained by a process that comprises contacting either:

(A) the modified aluminum oxy compound of claim 29,

(B) a transition metal compound, and

(C) an organoaluminum compound; or

(A) the modified aluminum oxy compound of claim 29,

U.S. Serial No. 10/734,218
Page 9

Amendment and Response to Restriction Requirement

(B) a transition metal compound,

(C) an organoaluminum compound, and

(D) any one of (D1) a boron compound represented by the general formula $BQ^1Q^2Q^3$, (D2) a boron compound represented by the general formula $G^+(BQ^1Q^2Q^3Q^4)^-$ and (D3) a boron compound represented by the general formula $(L-H)^+(BQ^1Q^2Q^3Q^4)^-$, wherein B represents a boron atom in the trivalent valence state; Q^1 to Q^3 may be the same or different and represent a halogen atom, a hydrocarbon group, a halogenated hydrocarbon group, a substituted silyl group, an alkoxy group or a di-substituted amino group; G^+ represents an inorganic or organic cation; and $(L-H)^+$ represents a Brønsted acid.

35. (Previously presented) A polymerization catalyst obtained by a process that comprises contacting either:

(A) the modified aluminum oxy compound of claim 30,

(B) a transition metal compound, and

(C) an organoaluminum compound; or

(A) the modified aluminum oxy compound of claim 30,

(B) a transition metal compound,

(C) an organoaluminum compound, and

(D) any one of (D1) a boron compound represented by the general formula $BQ^1Q^2Q^3$, (D2) a boron compound represented by the general formula $G^+(BQ^1Q^2Q^3Q^4)^-$ and (D3) a boron compound

U.S. Serial No. 10/734,218
Page 10

Amendment and Response to Restriction Requirement

represented by the general formula $(L-H)^+(BQ^1Q^2Q^3Q^4)^-$, wherein B represents a boron atom in the trivalent valence state; Q^1 to Q^3 may be the same or different and represent a halogen atom, a hydrocarbon group, a halogenated hydrocarbon group, a substituted silyl group, an alkoxy group or a di-substituted amino group; G^+ represents an inorganic or organic cation; and $(L-H)^+$ represents a Brønsted acid.

36. (Original) A process for producing an olefin polymer, which comprises homopolymerizing an olefin or copolymerizing olefins with the polymerization catalyst of claim 26.

37. (Original) A process for producing an olefin polymer, which comprises homopolymerizing an olefin or copolymerizing olefins with the polymerization catalyst of claim 27.

38. (Original) A process for producing an olefin polymer, which comprises homopolymerizing an olefin or copolymerizing olefins with the polymerization catalyst of claim 28.

39. (Original) A process for producing an olefin polymer, which comprises homopolymerizing an olefin or copolymerizing olefins with the polymerization catalyst of claim 29.

U.S. Serial No. 10/734,218
Page 11

Amendment and Response to Restriction Requirement

40. (Original) A process for producing an olefin polymer, which comprises homopolymerizing an olefin or copolymerizing olefins with the polymerization catalyst of claim 30.

41. (Original) A process for producing an olefin polymer, which comprises homopolymerizing an olefin or copolymerizing olefins with the polymerization catalyst of claim 31.

42. (Original) A process for producing an olefin polymer, which comprises homopolymerizing an olefin or copolymerizing olefins with the polymerization catalyst of claim 32.

43. (Original) A process for producing an olefin polymer, which comprises homopolymerizing an olefin or copolymerizing olefins with the polymerization catalyst of claim 33.

44. (Original) A process for producing an olefin polymer, which comprises homopolymerizing an olefin or copolymerizing olefins with the polymerization catalyst of claim 34.

45. (Original) A process for producing an olefin polymer, which comprises homopolymerizing an olefin or copolymerizing olefins with the polymerization catalyst of claim 35.

U.S. Serial No. 10/734,218
Page 12

Amendment and Response to Restriction Requirement

46. (Original) A process for producing an olefin polymer according to claim 36, wherein said olefin polymer is a copolymer of ethylene and α -olefin having 3 to 20 carbon atoms.

47. (Original) A process for producing an olefin polymer according to claim 37, wherein said olefin polymer is a copolymer of ethylene and α -olefin having 3 to 20 carbon atoms.

48. (Original) A process for producing an olefin polymer according to claim 38, wherein said olefin polymer is a copolymer of ethylene and α -olefin having 3 to 20 carbon atoms.

49. (Original) A process for producing an olefin polymer according to claim 39, wherein said olefin polymer is a copolymer of ethylene and α -olefin having 3 to 20 carbon atoms.

50. (Original) A process for producing an olefin polymer according to claim 40, wherein said olefin polymer is a copolymer of ethylene and α -olefin having 3 to 30 carbon atoms.

51. (Original) A process for producing an olefin polymer according to claim 36, wherein said olefin polymer is a homopolymer of 1-butene.

U.S. Serial No. 10/734,218
Page 13

Amendment and Response to Restriction Requirement

52. (Original) A process for producing an olefin polymer according to claim 37, wherein said olefin polymer is a homopolymer of 1-butene.

53. (Original) A process for producing an olefin polymer according to claim 38, wherein said olefin polymer is a homopolymer of 1-butene.

54. (Original) A process for producing an olefin polymer according to claim 39, wherein said olefin polymer is a homopolymer of 1-butene.

55. (Original) A process for producing an olefin polymer according to claim 40, wherein said olefin polymer is a homopolymer of 1-butene.

56. (Original) A process for producing an alkenyl aromatic hydrocarbon polymer, which comprises homopolymerizing an alkenyl aromatic hydrocarbon or copolymerizing at least one alkenyl aromatic hydrocarbon and at least one olefin with the polymerization catalyst of claim 26.

57. (Original) A process for producing an alkenyl aromatic hydrocarbon polymer, which comprises homopolymerizing an alkenyl aromatic hydrocarbon or copolymerizing at least one alkenyl aromatic hydrocarbon and at least one olefin with the polymerization catalyst of claim 27.

U.S. Serial No. 10/734,218
Page 14

Amendment and Response to Restriction Requirement

58. (Original) A process for producing an alkenyl aromatic hydrocarbon polymer, which comprises homopolymerizing an alkenyl aromatic hydrocarbon or copolymerizing at least one alkenyl aromatic hydrocarbon and at least one olefin with the polymerization catalyst of claim 28.

59. (Original) A process for producing an alkenyl aromatic hydrocarbon polymer, which comprises homopolymerizing an alkenyl aromatic hydrocarbon or copolymerizing at least one alkenyl aromatic hydrocarbon and at least one olefin with the polymerization catalyst of claim 29.

60. (Original) A process for producing an alkenyl aromatic hydrocarbon polymer, which comprises homopolymerizing an alkenyl aromatic hydrocarbon or copolymerizing at least one alkenyl aromatic hydrocarbon and at least one olefin with the polymerization catalyst of claim 30.

61. (Original) A copolymer of at least one alkenyl aromatic hydrocarbon and at least one olefin, having a number average molecular weight of 700,000 or more and a molecular weight distribution in terms of a ratio (M_w/M_n) of weight average molecular weight (M_w) to number average molecular weight (M_n), of 1.85 to 2.5.

U.S. Serial No. 10/734,218
Page 15

Amendment and Response to Restriction Requirement

62. (Original) An α -olefin polymerization catalyst according to claim 26, wherein the transition metal compound (B) is a transition metal compound having a capability of stereoregular polymerization of an α -olefin.

63. (Original) An α -olefin polymerization catalyst according to claim 27, wherein the transition metal compound (B) is a transition metal compound having a capability of stereoregular polymerization of an α -olefin.

64. (Original) An α -olefin polymerization catalyst according to claim 28, wherein the transition metal compound (B) is a transition metal compound having a capability of stereoregular polymerization of an α -olefin.


65. (Original) An α -olefin polymerization catalyst according to claim 29, wherein the transition metal compound (B) is a transition metal compound having a capability of stereoregular polymerization of an α -olefin.

66. (Original) An α -olefin polymerization catalyst according to claim 30, wherein the transition metal compound (B) is a transition metal compound having a capability of stereoregular polymerization of an α -olefin.


U.S. Serial No. 10/734,218
Page 16

Amendment and Response to Restriction Requirement

67. (Previously presented) An α -olefin polymerization catalyst according to claim 62, wherein the transition metal compound having a capability of stereoregular polymerization of an α -olefin is a transition metal compound (b1) represented by the general formula (4) or (5) below:



(wherein M is a transition metal atom of the Group IV of the Periodic Table, L^1 is an η^5 -indenyl group or a substituted η^5 -indenyl group, and two L^1 's may be mutually the same or different; Y^1 is a carbon atom, a silicon atom, a germanium atom or a tin atom, each of R^{11} and X^1 is a hydrogen atom, a halogen atom, an alkyl group, an aralkyl group, an aryl group, a substituted silyl group, an alkoxy group, an aralkyloxy group, an aryloxy group or a heterocyclic group, and all of R^{11} and X^1 may be the same or different mutually, P is 1 or 2); or




U.S. Serial No. 10/734,218
Page 17

Amendment and Response to Restriction Requirement

(wherein M is a transition metal atom of the Group IV of the Periodic Table, Y² is a silicon atom, a germanium atom or a tin atom, each of (R¹²_n-C₅H_{4-n}) and R¹²_q-C₅H_{4-q}) is a substituted η⁵-cyclopentadienyl group, and each of n and q is an integer of 1 to 3; the respective R¹² may be mutually the same or different, and indicate a halogen atom, an alkyl group, an aralkyl group, an aryl group, a substituted silyl group, an alkoxy group, an aralkyloxy group, an aryloxy group or a heterocyclic group.; the position and/or kind of R¹² in the substituted η⁵-cyclopentadienyl group is selected so that a symmetric plane including M does not exist; each of R¹³ and X² is a hydrogen atom, a halogen atom, an alkyl group, an aralkyl group, an aryl group, a substituted silyl group, an alkoxy group, an aralkyloxy group, an aryloxy group or a heterocyclic group, and all of R¹³ and X² may be the same or different mutually).


68. (Original) An α-olefin polymerization catalyst according to claim 63, wherein the transition metal compound having a capability of stereoregular polymerization of an α-olefin is a transition metal compound (b1) represented by the general formula (4) or (5) below:



U.S. Serial No. 10/734,218
Page 18

Amendment and Response to Restriction Requirement

(wherein M is a transition metal atom of the Group IV of the Periodic Table, L^1 is an η^5 -indenyl group or a substituted η^5 -indenyl group, and two L^1 's may be mutually the same or different; Y^1 is a carbon atom, a silicon atom, a germanium atom or a tin atom, each of R^{11} and X^1 is a hydrogen atom, a halogen atom, an alkyl group, an aralkyl group, an aryl group, a substituted silyl group, an alkoxy group, an aralkyloxy group, an aryloxy group or a heterocyclic group, and all of R^{11} and X^1 may be the same or different mutually, P is 1 or 2); or



(wherein M is a transition metal atom of the Group IV of the Periodic Table, Y^2 is a silicon atom, a germanium atom or a tin atom, each of $(R^{12}_n - C_5 H_{4-n})$ and $(R^{12}_q - C_5 H_{4-q})$ is a substituted η^5 -cyclopentadienyl group, and each of n and q is an integer of 1 to 3, the respective R^{12} may be mutually the same or different, and indicate a halogen atom, an alkyl group, an aralkyl group, an aryl group, a substituted silyl group, an alkoxy group, an aralkyloxy group, an aryloxy group or a heterocyclic group; the position and/or kind of R^{12} in the substituted η^5 -cyclopentadienyl group is selected so that a symmetric plane including M does not exist; each of R^{13} and X^2 is a hydrogen atom, a halogen atom, an alkyl group, an aralkyl group, an aryl group, a substituted silyl group, an

U.S. Serial No. 10/734,218
Page 19

Amendment and Response to Restriction Requirement

alkoxy group, an aralkyloxy group, an aryloxy group or a heterocyclic group, and all of R^{13} and X^2 may be the same or different mutually).


69. (Original) An α -olefin polymerization catalyst according to claim 64, wherein the transition metal compound having a capability of stereoregular polymerization of an α -olefin is a transition metal compound (b1) represented by the general formula (4) or (5) below:



(wherein M is a transition metal atom of the Group IV of the Periodic Table, L^1 is an η^5 -indenyl group or a substituted η^5 -indenyl group, and two L^1 's may be mutually the same or different; Y^1 is a carbon atom, a silicon atom, a germanium atom or a tin atom, each of R^{11} and X^1 is a hydrogen atom, a halogen atom, an alkyl group, an aralkyl group, an aryl group, a substituted silyl group, an alkoxy group, an aralkyloxy group, an aryloxy group or a heterocyclic group, and all of R^{11} and X^1 may be the same or different mutually, P is 1 or 2); or

U.S. Serial No. 10/734,218
Page 20

Amendment and Response to Restriction Requirement



(wherein M is a transition metal atom of the Group IV of the Periodic Table, Y² is a silicon atom, a germanium atom or a tin atom, each of (R¹²_n-C₅H_{4-n}) and (R¹²_q-C₅H_{4-q}) is a substituted η⁵-cyclopentadienyl group, and each of n and q is an integer of 1 to 3, the respective R¹² may be mutually the same or different, and indicate a halogen atom, an alkyl group, an aralkyl group, an aryl group, a substituted silyl group, an alkoxy group, an aralkyloxy group, an aryloxy group or a heterocyclic group, the position and/or kind of R¹² in the substituted η⁵-cyclopentadienyl group is selected so that a symmetric plane including M does not exist; each of R¹³ and X² is a hydrogen atom, a halogen atom, an alkyl group, an aralkyl group, an aryl group, a substituted silyl group, an alkoxy group, an aralkyloxy group, an aryloxy group or a heterocyclic group, and all of R¹³ and X² may be the same or different mutually).


70. (Original) An α-olefin polymerization catalyst according to claim 65, wherein the transition metal compound having a capability of stereoregular polymerization of an α-olefin is a transition metal compound (b1) represented by the general formula (4) or (5) below:

U.S. Serial No. 10/734,218
Page 21

Amendment and Response to Restriction Requirement



(wherein M is a transition metal atom of the Group IV of the Periodic Table, L^1 is an η^5 -indenyl group or a substituted η^5 -indenyl group, and two L^1 's may be mutually the same or different; Y^1 is a carbon atom, a silicon atom, a germanium atom or a tin atom, each of R^{11} and X^1 is a hydrogen atom, a halogen atom, an alkyl group, an aralkyl group, an aryl group, a substituted silyl group, an alkoxy group, an aralkyloxy group, an aryloxy group or a heterocyclic group, and all of R^{11} and X^1 may be the same or different mutually, P is 1 or 2); or




(wherein M is a transition metal atom of the Group IV of the Periodic Table, Y^2 is a silicon atom, a germanium atom or a tin atom, each of $(\text{R}^{12} \text{C}_5 \text{H}_{4-n})$ and $(\text{R}^{12} \text{C}_5 \text{H}_{4-q})$ is a substituted η^5 -cyclopentadienyl group, and each of n and q is an integer of 1 to 3, the respective R^{12} may be mutually the same or different, and indicate a halogen atom, an alkyl group, an aralkyl group, an aryl

U.S. Serial No. 10/734,218
Page 22

Amendment and Response to Restriction Requirement

group, a substituted silyl group, an alkoxy group, an aralkyloxy group, an aryloxy group or a heterocyclic group; the position and/or kind of R^{12} in the substituted η^5 -cyclopentadienyl group is selected so that a symmetric plane including M does not exist; each of R^{13} and X^2 is a hydrogen atom, a halogen atom, an alkyl group, an aralkyl group, an aryl group, a substituted silyl group, an alkoxy group, an aralkyloxy group, an aryloxy group or a heterocyclic group, and all of R^{13} and X^2 may be the same or different mutually).

71. (Original) An α -olefin polymerization catalyst according to claim 66, wherein the transition metal compound having a capability of stereoregular polymerization of an α -olefin is a transition metal compound (b1) represented by the general formula (4) or (5) below:




(wherein M is a transition metal atom of the Group IV of the Periodic Table, L^1 is an η^5 -indenyl group or a substituted η^5 -indenyl group, and two L^1 's may be mutually the same or different; Y^1

U.S. Serial No. 10/734,218
Page 23

Amendment and Response to Restriction Requirement

is a carbon atom, a silicon atom, a germanium atom or a tin atom, each of R^{11} and X^1 is a hydrogen atom, a halogen atom, an alkyl group, an aralkyl group, an aryl group, a substituted silyl group, an alkoxy group, an aralkyloxy group, an aryloxy group or a heterocyclic group, and all of R^{11} and X^1 may be the same or different mutually, P is 1 or 2); or



(wherein M is a transition metal atom of the Group IV of the Periodic Table, Y^2 is a silicon atom, a germanium atom or a tin atom, each of $(R^{12}_n - C_5 - H_{4-n})$ and $(R^{12}_q - C_5 - H_{4-q})$ is a substituted η^5 -cyclopentadienyl group, and each of n and q is an integer of 1 to 3; the respective R^{12} may be mutually the same or different, and indicate a halogen atom, an alkyl group, an aralkyl group, an aryl group, a substituted silyl group, an alkoxy group, an aralkyloxy group, an aryloxy group or a heterocyclic group; the position and/or kind of R^{12} in the substituted η^5 -cyclopentadienyl group is selected so that a symmetric plane including M does not exist; each of R^{13} and X^2 is a hydrogen

U.S. Serial No. 10/734,218
Page 24

Amendment and Response to Restriction Requirement

atom, a halogen atom, an alkyl group, an aralkyl group, an aryl group, a substituted silyl group, an alkoxy group, an aralkyloxy group, an aryloxy group or a heterocyclic group, and all of R^{13} and X^2 may be the same or different mutually).

72. (Original) A process for producing an α -olefin polymer, which comprises polymerizing an α -olefin with the α -olefin polymerization catalyst of claim 62.

73. (Original) A process for producing an α -olefin polymer, which comprises polymerizing an α -olefin with the α -olefin polymerization catalyst of claim 63.

74. (Original) A process for producing an α -olefin polymer, which comprises polymerizing an α -olefin with the α -olefin polymerization catalyst of claim 64.

75. (Original) A process for producing an α -olefin polymer, which comprises polymerizing an α -olefin with the α -olefin polymerization catalyst of claim 65.

76. (Original) A process for producing an α -olefin polymer, which comprises polymerizing an α -olefin with the α -olefin polymerization catalyst of claim 66.

U.S. Serial No. 10/734,218
Page 25

Amendment and Response to Restriction Requirement

77. (Original) The process for producing an α -olefin polymer according to claim 72, wherein the α -olefin polymer is an isotactic propylene polymer.

78. (Original) The process for producing an α -olefin polymer according to claim 73, wherein the α -olefin polymer is an isotactic propylene polymer.

79. (Original) The process for producing an α -olefin polymer according to claim 74, wherein the α -olefin polymer is an isotactic propylene polymer.

80. (Original) The process for producing an α -olefin polymer according to claim 75, wherein the α -olefin polymer is an isotactic propylene polymer.

81. (Original) The process for producing an α -olefin polymer according to claim 76, wherein the α -olefin polymer is an isotactic propylene polymer.